

Two cold-inducible genes encoding lipid transfer protein LTP4 from barley show differential responses to bacterial pathogens

Abstract The barley genes *HvLtp4.2* and *HvLtp4.3* both encode the lipid transfer protein LTP4 and are less than 1 kb apart in tail-to-tail orientation. They differ in their non-coding regions from each other and from the gene corresponding to a previously reported *Ltp4* cDNA (now *Ltp4.1*). Southern blot analysis indicated the existence of three or more *Ltp4* genes per haploid genome and showed considerable polymorphism among barley cultivars. We have investigated the transient expression of genes *HvLtp4.2* and *HvLtp4.3* following transformation by particle bombardment, using promoter fusions to the β -glucuronidase reporter sequence. In leaves, activities of the two promoters were of the same order as those of the sucrose synthase (*Ss1*) and cauliflower mosaic virus 35S promoters used as controls. Their expression patterns were similar, except that *Ltp4.2* was more active than *Ltp4.3* in endosperm, and *Ltp4.3* was active in roots, while *Ltp4.2* was not. The promoters of both genes were induced by low temperature, both in winter and spring barley cultivars. Northern blot analysis, using the *Ltp4*-specific probe, indicated that *Xanthomonas campestris* pv. *translucens* induced an increase over basal levels of *Ltp4* mRNA, while *Pseudomonas syringae* pv. *japonica* caused a decrease. The *Ltp4.3*-*Gus* promoter fusion also responded in opposite ways to these two compatible bacterial pathogens, whereas the *Ltp4.2*-*Gus* construction did not respond to infection.

Key words Bacterial pathogens · Barley · Lipid transfer proteins · Low temperature · Promoter analysis

Introduction

Non-specific lipid transfer proteins (LTPs) that stimulate the transfer of various lipids between membranes and artificial vesicles in vitro have been isolated from a wide range of plant sources (see Arondel and Kader 1990; Yamada 1992). A cytoplasmic role has been excluded for LTPs because they are generally secreted (Mundy and Rogers 1986; Sterk et al. 1991) and are associated with the cell wall (Molina et al. 1993; Molina and García-Olmedo 1993; Segura et al. 1993; Thoma et al. 1993; Pyee et al. 1994). Their possible involvement in the secretion or deposition of extracellular lipophilic molecules, such as cutin or wax, is under investigation (Sterk et al. 1991; Meijer et al. 1993; Hendriks et al. 1994; Pyee et al. 1994). Apart from their developmentally regulated expression, *Ltp* genes have been reported to respond to environmental stresses in tomato (Plant et al. 1991; Torres-Schumann et al. 1992; Kahn et al. 1993) and to cold treatment in barley (Dunn et al. 1991, 1994; Hughes et al. 1992; White et al. 1994). Evidence in support of a defense role for LTPs has been recently reviewed (García-Olmedo et al. 1995). These proteins are active against both bacterial and fungal pathogens in vitro (Molina and García-Olmedo 1991; Terras et al. 1992; Molina et al. 1993; Segura et al. 1993), and are distributed over exposed surfaces and in the vascular tissues, at basal concentrations that can be inhibitory for sensitive pathogens (Molina and García-Olmedo 1993; Pyee et al. 1994). Genes encoding these proteins respond to bacterial and fungal infections in a complex manner (Molina and García-Olmedo 1994; García-Olmedo et al. 1995, 1996). Furthermore, transgenic plants expressing LTP show enhanced tolerance to a variety of pathogens (Molina et al., in preparation) and LTP-sensitive mutants of bacterial pathogens have been shown to be avirulent (Titarenko et al., in preparation).

In barley, one aleurone-specific and four ubiquitous LTPs have been purified (Mundy and Rogers 1986; Molina et al. 1993), and several of the corresponding

cDNAs and genes have been cloned (Linnestead et al. 1991; Skriver et al. 1992; Dunn et al. 1991; Molina and García-Olmedo 1993; Gausing 1994; White et al. 1994). Leaf-expressed *Ltp* genes have been shown to be induced by pathogens (Molina and García-Olmedo 1993; García-Olmedo et al. 1995) and although a cold response was observed in the winter barley cv. Igri (Dunn et al. 1991; Hughes et al. 1992; White et al. 1994), such a response was not observed in spring barley cv. Bomi (Molina and García-Olmedo 1993).

We have now characterized two *Ltp* genes, *HvLtp4.2* and *HvLtp4.3*, both of which encode the same mature protein as a previously reported LTP4 cDNA (Molina and García-Olmedo 1993), whose corresponding gene (now *HvLtp4.1*; syn. *gblt4.2* from White et al. 1994) is different from *HvLtp4.2* and *HvLtp4.3*. Gene *HvLtp4.2* is practically identical to that in clone *gblt4.9* (White et al. 1994), whereas *HvLtp4.3* is significantly divergent. We have investigated the activity of these two promoters, using fusions to the β -glucuronidase reporter (GUS) gene and particle bombardment, and have found that while both promoters are induced by cold treatment, only one of them (*HvLtp4.3*) responds to pathogen infection.

Materials and methods

Cloning of *HvLtp4.2* and *HvLtp4.3*

Lambda EMBL3 recombinant clones (1.2×10^6) of a genomic library from *Hordeum vulgare* L. var. NK1558 (spring habit; Clontech), were transferred to Hybond N membranes (Amersham) and hybridized at 65°C with the cDNA probe (Molina and García-Olmedo 1993) following standard procedures (Sambrook et al. 1989). Two of the selected clones (*λltp4.11* and *λltp4.14*) were purified through two further rounds of screening and restriction mapped. The *Bam*HI fragments hybridizing with the probe were subcloned into pBluescript II KS (Stratagene) and sequenced according to Hattoni and Sakaki (1986).

DNA and RNA hybridization

DNA was isolated essentially as described (Taylor and Powell 1982), restricted with *Bam*HI endonuclease, subjected to electrophoresis in 0.8% agarose, and transferred to Hybond N membranes (Amersham) following standard procedures. Hybridization at 65°C was according to Church and Gilbert (1984).

RNA was purified from frozen tissues by phenol/chloroform extraction, followed by precipitation with 3 M lithium chloride (Lagrimini et al. 1987), and subjected to electrophoresis (7.5 µg each) in 7.5% formaldehyde/agarose gels, which were blotted to Hybond N membranes (Amersham). Hybridization was carried out at 65°C as above. Ethidium bromide (40 µg/ml) was included in the loading buffer to allow visualization under UV light and photography. Equal sample loads were checked by densitometry (INH Image programme, Bio-Rad) of the negatives of the UV photographs. Quantitation of radioactive signals in Northern-blot experiments was carried out by densitometry (INH Image programme, Bio-Rad) of the autoradiographs from three independent filters.

Gene constructions and particle bombardment

Fragments corresponding to the promoters of *HvLtp4.2* (positions -1 to -665 nt) and *HvLtp4.3* (-1 to -755 nt) were fused in-phase to the β -glucuronidase reporter gene (Jefferson 1987) in the pBluescript II KS (Stratagene) vector (Fig. 1A). The indicated region of *Ltp4.2* promoter was amplified by the polymerase chain reaction from a *Bam*HI subclone of the gene (Fig. 1A) and the equivalent region (as determined by sequence alignment) from promoter *Ltp4.3* was similarly amplified from a *Sall*-*Eco*RI subclone of the corresponding gene (Fig. 1A). Gene constructions used as controls have been previously described: 35S-GUS contained the coding region of the *Gus* (*uidA*) gene and the 35S cauliflower mosaic virus (CaMV) promoter, subcloned in the pUC19 vector (Diaz and Carbonero 1992); 35Sf-GUS additionally contained intron 1 of the maize *Adh1* gene (Vasil et al. 1991); *Ubi*-GUS and *Ubi*-LUC contained the maize ubiquitin promoter-exon 1-intron 1, fused either to the GUS or the luciferase reporter genes (Christensen et al. 1992); *lir1*-GUS carried the endosperm-specific promoter of the barley trypsin inhibitor (Diaz et al. 1993); in *Ss1*-GUS the reporter gene is controlled by the promoter and first intron of barley sucrose synthase gene *Ss1* (Diaz et al. 1995).

Particle bombardment was carried out with a biolistic helium gun device (DuPont PDS-100 λ). Gold particles (1.0 µm in size) were coated with DNA as described by Taylor and Vasil (1991). Barley tissues were placed onto a Phytigel-H₂O support (Sigma) in 6-cm diameter dishes and each plate was bombarded once with 0.154 µg gold and 0.350 µg DNA at a distance of 7.5 cm from the macro-projectile stopper shelf. After bombardment, dishes were incubated at 25°C for 2 days. Standard fluorometric GUS assays were performed using 4-methyl-umbelliferyl- β -D-glucuronide (MUG) as substrate (Jefferson 1987) and were quantified with a Heraeus TK 100 fluorimeter. Luciferase assays were done as described by Callis et al. (1987), using a 15-s integration time on a luminometer (Analytical Luminiscence Laboratory, Monolight 2010). Treatment with abscisic acid (10 µM) was carried out by including this hormone in the solid medium where leaves were bombarded and incubated for 24 h.


Infection with bacteria

Five-day-old barley (cv. Bomi) plants grown on vermiculite (22°C day/18°C night; 16 h light) were inoculated by infiltration with the bacteria *Pseudomonas syringae* pv. *japonica*, *Xanthomonas campestris* pv. *translucens* (10 µl of 10^8 cfu/ml in 10 mM MgCl₂), or with 10 mM MgCl₂ (mock inoculation). After inoculation, leaf samples were taken at times indicated in Fig. 5A and immediately frozen in liquid nitrogen. In the experiments with excised leaves, inoculations were carried out as above just before excision.

Results

Gene, *HvLtp4.2* and *HvLtp4.3*

Two genomic clones, *λltp4.11* and *λltp4.14*, which hybridized with a probe that consisted of the 3' untranslated region from the previously reported *Ltp4* cDNA (now *Ltp4.1*; Molina and García-Olmedo 1993), were isolated from a genomic barley library. As shown in Fig. 1A, both clones contained the same two *Ltp* genes, *HvLtp4.2* and *HvLtp4.3*, less than 1 kb apart in a tail-to-tail orientation. Their deduced mature protein sequences were identical and differed from that of the LTP4.1 in a conservative amino acid change (A/V), but



nucleotide sequences encoding the signal peptide and the 3' untranslated regions were distinct from each other. In Fig. 1B, an alignment of the two promoters shows that they are also divergent, specially with respect to key motifs potentially involved in their responses to pathogen infection.

Southern blot analysis with the *Ltp4*-specific probe, which recognized all three *Ltp4* genes, but not the other known *Ltp* genes, detected polymorphic patterns that were consistent with the existence of multiple copies per haploid genome (Fig. 2). Previously reported Northern blot analyses with the same probe must therefore have detected the overall expression of all the *Ltp4* genes (Molina and García-Olmedo 1993).

Gene fusions involving the *HvLtp4.2* and *HvLtp4.3* promoters and the GUS reporter (Fig. 1A) were transiently expressed following particle bombardment in barley leaves and activity levels were similar in both cases, and of the same order as those obtained with the barley sucrose synthase *Ss1* and CaMV 35S promoters used as positive controls (Fig. 3A). Under the same conditions, no significant activity was driven by the endosperm-specific promoter from barley *ltp1* gene, which was used as a negative control. Activities of the two *Ltp* gene constructions found in different tissues, using the *Ubi-LUC* chimeric fusion as internal control, were similar to each other and essentially reproduced the expression pattern obtained by Northern blot analysis, except that

Fig. 1 A Schematic representation of genomic clones containing genes *HvLtp4.2* and *HvLtp4.3* and of fusions of their promoters (P4.2 and P4.3) to the β -glucuronidase/nopaline synthase (GUS/NOS) reporter gene. Restriction endonuclease sites for *Bam*HI (B) and *Sal*I (S) are indicated. **B** Alignment of the 5' untranslated sequences of the two *Ltp* genes used in the gene fusions. Numbering is from the ATG translation initiation site; identities and gaps are indicated by asterisks and dashes, respectively. The following motifs have been highlighted: G-box, consensus ACGT (*Gbox*; Guiltinan et al. 1990; Schindler et al. 1992); H-box, consensus (C/A)A(C/A)C(T/A)A(C/A)C(*Hbox*; Lois et al. 1989; Cramer et al. 1989; Ohl et al. 1990; Loake et al. 1992; Yu et al. 1993); and AP-1, consensus TGACACA (*API*; Després et al. 1995). The alignment was constructed out using the CLUSTAL W programme (Thompson et al. 1994), and the SIGNAL SCAN 4.0 programme (Prestridge 1991) was used to search for sequence motifs

Ltp4.2 was more active than *Ltp4.3* in endosperm, and *Ltp4.3* was active in roots, while *Ltp4.2* was not (Fig. 3B).

Responses to low temperature

Experimental conditions were optimized using the *Ss1*-GUS gene fusion, as the sucrose synthase *Ss1* promoter had been previously shown to be cold responsive (Marañón et al. 1990). It was established that an induction




Fig. 5 A Responses of *Ltp4* genes to the indicated bacterial pathogens. Northern blot signals were obtained with the *Ltp4*-specific probe (Molina and García-Olmedo 1993) and quantitated by densitometry (INH Image programme, Bio-Rad). B Responses of promoter fusions *Ltp4.2* and *Ltp4.3* to the same pathogens. Groups of three leaves were inoculated (i) in the apical half leaf (a) and bombarded both in the apical and in the basal (b) parts of the leaf, using a plastic window in the positions indicated by a discontinuous line. The experiment was repeated three times for each permutation (1–3) of the treatment. GUS activity was determined after 24 h incubation in the Phytigel-H₂O support. The control (mock) value of the apical part was taken as 1 in each experiment. Standard errors of the mean are indicated by bars

at a basal site (non-inoculated half-leaves), as illustrated in Fig. 5B, and GUS activity was determined fluorometrically for each bombarded half-leaf. The two *Ltp* promoters responded in a differential manner to the pathogens: whereas the activity of promoter *Ltp4.2* was not affected by the bacterial infections, that of promoter *Ltp4.3* reproduced the responses detected by Northern analysis both in the apical and in the basal half of the leaf: an increase in response to infection by *X. campestris* and a decrease in the case of *P. syringae* (Fig. 5B). These results implied that the response of the

Ltp4.3 promoter to pathogens occurs beyond the infected zone of the leaf.

Discussion

A common characteristic of related genes is that their coding regions have diverged less than their non-coding regions. Genes *HvLtp4.2* and *HvLtp4.3* represent an extreme example, as they encode the same mature protein while their non-coding regions are quite different. However, this divergence does not seem simply to reflect less stringent selective constraints affecting these parts of the genes but to have positive functional relevance, as it has led to alterations in their expression patterns. The above results indicate that both genes characterized here are active (not pseudogenes), and confirmed the previously reported complexity of the *Ltp* gene family (Molina and García-Olmedo 1993; White et al. 1994). Their developmental expression patterns, as judged from the transient expression experiments, were similar – except for minor differences – and in general agreement with the overall LTP4 expression pattern previously observed by Northern blot analysis (Molina and García-Olmedo 1993).

The excised-leaf procedure that has been developed here to study responses to low temperature and to pathogen infection in barley, using the well-established particle bombardment technique, represents a simplification of previous whole-plant approaches and allows the rapid investigation of different promoters in multiple genetic backgrounds. Thus, it has been shown that the two *Ltp* promoters, which have been isolated from a spring barley, are induced by low temperature both in winter (cv. Igri) and in spring (cvs. Bomi and Himalaya) barleys, indicating that the observed response must depend on the promoter characteristics, and is not dependent on the genetic background of the cultivars used in the study or on their spring or winter habit. We have previously observed a non-significant cold-induced increase (<20%) in *Ltp4* mRNA levels in cv. Bomi by Northern blot analysis, under conditions in which the *Ss1* promoter underwent a 20-fold induction (Molina and García-Olmedo 1993), while an induction of *Ltp* genes by low temperature has been reported for cv. Igri (Dunn et al. 1991; Hughes et al. 1992; White et al. 1994). In view of the current results, the difference can be ascribed either to divergence in the *Ltp4* promoters of the two cultivars or to the presence of a highly expressed, non-responsive *Ltp4* gene in cv. Bomi that would mask the induction of other *Ltp4* genes. In contrast, a significant response of *Ltp4* genes to abscisic acid was detected both in winter and spring barley (Hughes et al. 1992; Molina and García-Olmedo 1993).

It has previously been shown that expression of *Ltp* genes can be altered, independently from each other

and from other defense genes, in different plant-pathogen interactions (Molina and García-Olmedo 1993, 1994; García-Olmedo et al. 1995, 1996). We have observed opposite responses in two compatible bacterium-plant interactions by Northern blot analysis, which detects the overall expression of *Ltp4* genes. The observed negative response to *P. syringae* infection has some precedents in the literature, including those of several defense genes in bean (Jakobek et al. 1993) and the *StPth1* gene in potato (Moreno et al. 1994), that have been interpreted as a mechanism by which some pathogens might overcome constitutive defense barriers.

The two *Ltp* promoters analysed here behave differentially with respect to pathogens: responses of promoter *Ltp4.3* mimic the overall *Ltp4* pattern observed by Northern blot analysis, while the activity of promoter *Ltp4.2* is not affected by either pathogen. This is consistent with the overall divergence of these promoters and, in particular, with the observed differences in putative pathogen-response motifs (Fig. 1B). The 5' untranslated regions of the two *Ltp* genes each contain a copy of an ACGT motif common to several cis-acting elements that bind basic leucine zipper proteins, including those denominated G-box (Schindler et al. 1992) and ABRE (Guiltinan et al. 1990). Perhaps of greater relevance is the presence in gene *HvLtp4.3* of the pathogen-responsive motifs H-box and AP-1 (Fig. 1B). The first motif is present in promoters of phenylpropanoid biosynthetic genes (Cramer et al. 1989; Lois et al. 1989; Ohl et al. 1990; Loake et al. 1992; Yu et al. 1993), which have been shown to display elicitor-inducible and light-inducible footprints in vivo (Lois et al. 1989). The presence of H-box and G-box motifs in the promoter of these genes has been found to be required for elicitor induction (Loake et al. 1992). The AP-1 element (TGACACA) has been described in the pathogenesis-related *PR-10a* gene from potato (Després et al. 1995), where it is involved, together with an E-box (CANNTG), in gene activation by an elicitor of *Phytophthora infestans* and interacts with nuclear factor PBF-1, which requires phosphorylation for activity (Després et al. 1995). Several E-box cores can be identified in the *HvLtp4.3* promoter (not highlighted in Fig. 1B). The variation in pathogen-response motifs among *Ltp* promoters is in agreement with the previously observed plasticity of defense-gene elicitation in plant-pathogen interactions and with the combinatorial nature of plant defense responses (Molina and García-Olmedo 1994; García-Olmedo et al. 1995, 1996).

Acknowledgements Sequencing of DNA by R. Novitzky (CIBA-GEIGY, Research Triangle Park, NC, USA) and technical assistance from D. Lamóneda and J. García are gratefully acknowledged. This work was financed by the DGICYT, Spain (Grants PB92-0325 and PB94-0404).

References

- Aronel V, Kader J-C (1990) Lipid transfer proteins in plants. *Experientia* 46: 579-585
- Callis J, Fromm M, Walbot W (1987) Introns increase gene expression in cultured maize cells. *Genes Dev* 1: 1183-1200
- Christensen AH, Sharrock RA, Quail PH (1992) Maize polyubiquitin genes: structure, thermal perturbation of expression and transcript splicing, and promoter activity following transfer to protoplasts by electroporation. *Plant Mol Biol* 18: 675-689
- Church GM, Gilbert W (1984) Genomic sequencing. *Proc Natl Acad Sci USA* 81: 1991-1995
- Cramer CL, Edwards K, Dron M, Liang X, Dildine SL, Bolwell GP, Dixon RA, Lamb CJ (1989) Phenylalanine ammonia-lyase gene organization and structure. *Plant Mol Biol* 12: 367-383
- Després C, Subramaniam R, Matton DP, Brisson N (1995) The activation of the potato *PR-10a* gene requires the phosphorylation of the nuclear factor PBF-1. *Plant Cell* 7: 589-598
- Díaz I, Carbonero P (1992) Isolation of protoplasts from developing barley endosperm: a tool for transient expression studies. *Plant Cell Rep* 10: 595-598
- Díaz I, Royo J, Carbonero P (1993) The promoter of barley trypsin-inhibitor BTI-CMe, discriminates between wheat and barley endosperm protoplasts in transient expression assays. *Plant Cell Rep* 12: 698-701
- Díaz I, Royo J, Sánchez de la Hoz P, Carbonero P (1995) Gene specificity is maintained in transient expression assays with protoplasts derived from different tissues of barley. *Euphytica* 85: 203-207
- Dunn MA, Hughes MA, Zhang L, Pearce RS, Quigley AS, Jack PL (1991) Nucleotide sequence and molecular analysis of the low temperature induced cereal gene, BLT4. *Mol Gen Genet* 229: 389-394
- Dunn MA, Goddard NJ, Zhang L, Pearce RS, Hughes MA (1994) Low-temperature-responsive genes have different control mechanisms. *Plant Mol Biol* 24: 879-888
- García-Olmedo F, Molina A, Segura A, Moreno M (1995) The defensive role of nonspecific lipid-transfer proteins in plants. *Trends Microbiol* 3: 72-74
- García-Olmedo F, Molina A, Segura A, Moreno M, Castagnaro A, Titarenko E, Rodríguez-Palenzuela P, Piñeiro M, Díaz I (1996) Engineering plants against pathogens: a general strategy. *Field Crop Res*, in press
- Gausling K (1994) Lipid transfer protein genes specifically expressed in barley leaves and coleoptiles. *Planta* 192: 574-580
- Guiltinan MJ, Marcotte WR, Quatrano RS (1990) A plant leucine zipper protein that recognizes an abscisic acid response element. *Science* 250: 267-271
- Hattoni M, Sakaki Y (1986) Dideoxy sequencing method using denatured plasmid templates. *Anal Biochem* 152: 232-238
- Hendriks T, Meijer EA, Thoma S, Kader J-C, de Vries SC (1994) The carrot extracellular lipid transfer protein EP2: quantitative aspects with respect to its putative role in cutin synthesis. In: Coruzzi G, Puigdomènech P (eds) *Plant Molecular Biology: NATO ASI Series vol H81*. Springer-Verlag, Berlin-Heidelberg, pp 85-94
- Hughes MA, Dunn MA, Pearce RS, White AJ, Zhang L (1992) An abscisic-acid-responsive, low temperature barley gene has homology with a maize phospholipid transfer protein. *Plant Cell Environ* 15: 861-865
- Jakobek JL, Smith JA, Lindgren PB (1993) Suppression of bean defense responses by *Pseudomonas syringae*. *Plant Cell* 5: 57-63
- Jefferson RA (1987) Assaying chimeric genes in plants: the GUS gene fusion system. *Plant Mol Biol Rep* 5: 387-405
- Kahn TL, Fender SE, Bray EA, O'Connell MA (1993) Characterization of expression of drought and abscisic acid-regulated tomato genes in the drought-resistant species *Lycopersicon pennellii*. *Plant Physiol* 103: 597-605

- Lagrimini LM, Burkhart W, Moyer M, Rothstein S (1987) Molecular cloning of complementary DNA encoding the lignin-forming peroxidase from tobacco: molecular analysis and tissue-specific expression. *Proc Natl Acad Sci USA* 84: 7542-7546
- Linnestad C, Lönneborg A, Kalla R, Olsen O-A (1991) Promoter of a lipid transfer protein gene expressed in barley aleurone cells contains similar *myb* and *myc* recognition sites as the maize *Bz-McC* allele. *Plant Physiol* 97: 841-843
- Loake GJ, Faktor O, Lamb CJ, Dixon RA (1992) Combination of H-box [CCTACC(N)₂CT] and G-box (CACGTG) cis elements is necessary for feed-forward stimulation of a chalcone synthase promoter by the phenylpropanoid-pathway intermediate *p*-coumaric acid. *Proc Natl Acad Sci USA* 89: 9230-9234
- Lois R, Dietrich A, Hahlbrock K, Schulz W (1989) A phenylalanine ammonia-lyase gene from parsley: structure, regulation and identification of elicitor and light responsive cis-acting elements. *EMBO J* 8: 1641-1648
- Maraña C, García-Olmedo F, Carbonero P (1990) Differential expression of two types of sucrose synthase-encoding genes in wheat in response to anaerobiosis, cold shock and light. *Gene* 88: 167-172
- Meijer EA, de Vries SC, Sterk P, Gadella DWJ, Wirtz KWA, Hendriks T (1993) Characterization of the non-specific lipid transfer protein EP2 from carrot (*Daucus carota* L.). *Mol Cell Biochem* 123: 159-166
- Molina A, García-Olmedo F (1991) Patent application P9101258 (24.05.91) PCT/EP92/01130. US Patent Number 5,446,127 (29.08.95).
- Molina A, García-Olmedo F (1993) Developmental and pathogen-induced expression of three barley genes encoding lipid transfer proteins. *Plant J* 4: 983-991
- Molina A, García-Olmedo F (1994) Expression of genes encoding thionins and lipid-transfer proteins. A combinatorial model for the responses of defense genes to pathogens. In: Coruzzi G, Puigdomènech P (eds) *Plant Molecular Biology: NATO ASI Series vol H81*. Springer-Verlag, Berlin-Heidelberg, pp 235-244
- Molina A, Segura A, García-Olmedo F (1993) Lipid transfer proteins (nsLTPs) from barley and maize leaves are potent inhibitors of bacterial and fungal plant pathogens. *FEBS Lett* 316: 119-122
- Moreno A, Segura A, García-Olmedo F (1994) Pseudothionin-St1, a potato peptide active against potato pathogens. *Eur J Biochem* 223: 135-139
- Mundy J, Rogers JC (1986) Selective expression of a probable amylase/protease inhibitor in barley aleurone cells: comparison to the barley amylase/subtilisin inhibitor. *Planta* 169: 51-63
- Ohl S, Hedrick SA, Chory J, Lamb CJ (1990) Functional properties of a phenylalanine ammonia-lyase promoter from *Arabidopsis*. *Plant Cell* 2: 837-848
- Plant AL, Cohen A, Moses MS, Bray EA (1991) Nucleotide sequence and spatial expression pattern of a drought- and abscisic acid-induced gene of tomato. *Plant Physiol* 97: 900-906
- Prestridge DS (1991) SIGNAL SCAN: a computer program that scans DNA sequences for eukaryotic transcriptional elements. *CABIOS* 7: 203-206
- Pyee J, Yu H, Kolattukudy PE (1994) Identification of a lipid transfer protein as the major protein in the surface wax of broccoli (*Brassica oleracea*) leaves. *Arch Biochem Biophys* 311: 460-468
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual* (2nd edn). Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York
- Schindler V, Menkens AE, Beckmann H, Ecker JR, Cashmore AR (1992) Heterodimerization between light-regulated and ubiquitously expressed *Arabidopsis* GBF bZIP proteins. *EMBO J* 11: 1261-1273
- Segura A, Moreno M, García-Olmedo F (1993) Purification and antipathogenic activity of lipid transfer proteins (LTPs) from leaves of *Arabidopsis* and spinach. *FEBS Lett* 332: 243-246
- Skriver K, Leah R, Müller-Urli F, Olsen F-L, Mundy J (1992) Structure and expression of the barley lipid transfer protein gene *Ltp1*. *Plant Mol Biol* 18: 585-589
- Sterk P, Booiij H, Schellekens GA, Van Kammen A, de Vries SC (1991) Cell-specific expression of the carrot EP2 lipid transfer protein gene. *Plant Cell* 3: 907-921
- Taylor B, Powell A (1982) Isolation of plant DNA and RNA. *Focus* 4: 4-5
- Taylor MG, Vasil IK (1991) Histology of, and physical factors affecting, transient GUS expression in pearl millet (*Pennisetum glaucum* (L.) R.Br.) embryos following microprojectile bombardment. *Plant Cell Rep* 10: 120-125
- Terras FRG, Goderis IJ, Van Leuven F, Vanderleyden J, Cammue BPA, Broekaert WF (1992) In vitro antifungal activity of a radish (*Raphanus sativus* L.) seed protein homologous to nonspecific lipid transfer proteins. *Plant Physiol* 100: 1055-1058
- Thoma S, Kaneko Y, Somerville C (1993) A non-specific lipid transfer protein from *Arabidopsis* is a cell wall protein. *Plant J* 3: 427-436
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673-4680
- Torres-Schumann S, Godoy JA, Pintor-Toro JA (1992) A probable lipid transfer protein gene is induced by NaCl in stems of tomato plants. *Plant Mol Biol* 18: 749-757
- Vasil V, Brown SM, Re D, Fromm ME, Vasil IK (1991) Stably transformed callus lines from microprojectile bombardment of cell suspension cultures of wheat. *Biotechnology* 9: 743-746
- White AJ, Dunn MA, Brown K, Hughes MA (1994) Comparative analysis of genomic sequences and expression of a lipid transfer protein gene family in winter barley. *J Exp Botany* 45: 1885-1892
- Yamada M (1992) Lipid transfer proteins in plants and microorganisms. *Plant Cell Physiol* 33: 1-6
- Yu LM, Lamb CJ, Dixon RA (1993) Purification and biochemical characterization of proteins which bind to the H-box cis-element implicated in transcriptional activation of plant defense genes. *Plant J* 3: 805-816